


IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

VERIFIED DECLARATION

I, ROGER CHARLES STANFORD TUNN, Technical Translator of 4, Plymouth Park, Sevenoaks, Kent, England, do hereby declare that I am conversant with the German and English languages and am a competent translator thereof. I declare further that my name is included in the list of people from whom the Comptroller is prepared to accept translations from the German language and that the attached is a true translation into the English language made by me of an official copy of the document in respect of a patent application filed in Germany on 30th December 1997 under No. 197 58 178.1 and of the official certificate attached thereto.

I further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

I hereby subscribe my name to the foregoing declaration, this 13th day of October, 2003.



Roger Charles Stanford Tunn

TECHNOLOGY CENTER 1700

OCT 27 2003

RECEIVED

THIS PAGE BLANK (USPTO)

Translation:

FEDERAL REPUBLIC OF GERMANY

Priority Certificate in Respect of the Filing of a Patent Application

Serial No.: 197 58 178.1

Filing date: 30th December 1997

Applicant: Henkel Kommanditgesellschaft auf Aktien, Düsseldorf/DE

Title: Dishwasher detergent shaped bodies containing chlorine
bleaching agents

IPC: C 11 D 17/00

The attached documents are a true and exact reproduction of the
original documents of that patent application.

(SEAL)

Munich, 24th September, 2003

German Patent and Trade Mark Office

The President:

Per Pro
(signature)

RECEIVED
OCT 27 2003
TECHNOLOGY CENTER 1700

THIS PAGE BLANK (USPTO)

Henkel KGaA
Dr. Pressner
30.12.1997

Patent Application H 3174

Dishwasher Detergent Shaped Bodies Containing Chlorine Bleaching
Agents

5 This invention relates to deterative shaped bodies, primarily tablets, such as dishwasher tablets, laundry detergent tablets, bleach tablets, stain remover tablets or water softening tablets, for use in the home, more particularly for use in machines, to a process for their production and to their use.

Deterative shaped bodies, more particularly tablets, have a number of advantages over powder-form compositions, such as easy handling, simple dosage and low packaging volumes.

10 Unfortunately, problems arise out of the fact that, to achieve adequate dimensional stability and fracture resistance, relatively high pressures have to be applied in the compression of the powder-form ingredients. In view of the heavy compaction which they undergo, corresponding tablets often show inadequate storage stability in regard to sensitive ingredients (bleaching agent, perfume, enzymes, silver
15 protectors, dyes, surfactants), reduced performance (particularly against such soils as tea, milk, starch) by comparison with comparable amounts of active ingredient made up in another form, occasionally inadequate protection of sensitive materials to be cleaned and impaired disintegration and dissolving properties in use.

20 According to some patents belonging to the prior art, some of these problems are said to have been solved.

In many cases, however, the consumer is unable to reproduce the performance advantages described in those documents. Accordingly, an

important problem confronting manufacturers of deterative shaped bodies is to satisfy the consumer.

Accordingly, the problem addressed by the present invention was to provide high-performance, easy-to-handle deterative shaped bodies, above
5 all dishwasher tablets, laundry detergent tablets, stain remover tablets or water softener tablets for use in the home, more particularly for use in machines, with high storage stability, minimal packaging volume, a good fragrance profile and a defined solubility profile which, above all, would guarantee high consumer satisfaction.

10 This problem has surprisingly been solved by a shaped body in which an ingredient (I) is mainly located in a defined region, this region representing no more than 40% by volume of the shaped body.

Accordingly, the present invention relates to a shaped body containing builders, alkali sources, bleaching agents, enzymes and
15 surfactants, characterized in that more than 80% by weight of a mixture of active substance (I) in the form of a bleaching agent from the group of chlorine bleaching agents and a fusible component for controlling solubility selected from the group of surfactants, paraffins, microwaxes and relatively high molecular weight polyethylene glycols is present in a region of no
20 greater than 40% by volume.

Besides the bleaching agent from the group of chlorine bleaching agents, the ingredient (I) may also be a bleach activator, a silver protector and/or a soil-release compound, an enzyme, a surfactant or a component
25 or a compound for controlling solubility. However, the ingredient (I) may also be a mixture of these components.

In one particularly preferred embodiment, the ingredient (I) is a mixture of a component or a compound for controlling solubility and at least one other ingredient from the group consisting of bleaching agent and/or bleach activator and/or silver protector and/or soil-release compound
30 and/or enzyme and/or a surfactant.

The solubility of the region containing the ingredient (I), the surface, the nature of the compression and storage stability can also have a critical bearing on the properties of the tablet.

According to the invention, it is favorable if the region containing
5 more than 80% by weight, preferably more than 90% by weight, more preferably more than 95% by weight and, most preferably, 100% by weight of an ingredient (I) present is surrounded in exactly three directions in space by other ingredients of the shaped body.

In another preferred embodiment of the invention, at least one
10 region containing more than 80% by weight, preferably more than 90% by weight, more preferably more than 95% by weight and most preferably 100% by weight of an ingredient (I) present dissolves more than 5%, preferably more than 10%, more preferably more than 25%, most preferably more than 50% and, in one particularly advantageous
15 embodiment, more than 100% faster than the remaining region and/or regions of the shaped body.

According to the invention, it is also positive if more than 80% by weight, preferably more than 90% by weight, more preferably more than 95% by weight and most preferably 100% by weight of an ingredient (I)
20 present is contained in a region of no greater than 40% by volume, preferably between 5 and 30% by volume, more preferably between 10 and 25% by volume and most preferably between 15 and 20% by volume of the shaped body.

The overall result can also be desirably influenced if the increase in
25 weight of the region containing ingredient (I) during storage under normal domestic conditions, i.e. between 15 and 30°C and between 5 and 55% and preferably between 15 and 35% air humidity for a period of 30 days, preferably 60 days and more preferably 90 days is no more than 50% by weight, preferably no more than 40% by weight, more preferably no more
30 than 30% by weight, most preferably no more than 20% by weight and, in

one particularly advantageous embodiment, no more than 10% by weight and, in the most favorable case, no more than 5% by weight.

The overall result can also be desirably influenced if the loss of active substance (I) of the region during storage under normal domestic conditions, i.e. between 15 and 30°C and between 5 and 55% and preferably between 15 and 35% air humidity for a period of 30 days, preferably 60 days and more preferably 90 days is no more than 50% by weight, preferably no more than 40% by weight, more preferably no more than 30% by weight, most preferably no more than 20% by weight and, in one particularly advantageous embodiment, no more than 10% by weight and, in the most favorable case, no more than 5% by weight.

The overall result can also be desirably influenced if the absorption maximum of a 1% solution of a colored region or a colored component during storage under normal domestic conditions, i.e. between 15 and 30°C and between 5 and 55% and preferably between 15 and 35% air humidity for a period of 30 days, preferably 60 days and more preferably 90 days is no more than 100 wave numbers, preferably no more than 50 wave numbers, more preferably no more than 30 wave numbers, most preferably no more than 20 wave numbers and, in one particularly advantageous embodiment, no more than 10 wave numbers and, in the most favorable case, no more than 5 wave numbers.

The shaped body according to the invention is also positively influenced if the region containing more than 80% by weight, preferably more than 90% by weight, more preferably more than 95% by weight and, most preferably, 100% by weight of an ingredient (I) present is surrounded in exactly three directions in space by other ingredients of the shaped body.

Solubility

The solubility can be influenced by components and/or compounds

for accelerating solubility (disintegrators) or retarding solubility.

Suitable disintegrators are any of the disintegrators known from the prior art. Particular reference is made in this regard to *Römpp (9th Edition, Vol. 6, page 4440)* and *Voigt "Lehrbuch der pharmazeutischen Technologie" (6th Edition, 1987)*. Particularly suitable disintegrators are such materials as starch, cellulose and cellulose derivatives, alginates, dextrans, crosslinked polyvinyl pyrrolidones and others; systems of weak acids and carbonate-containing disintegrators, more particularly citric acid and tartaric acid in combination with hydrogen carbonate or carbonate and polyethylene glycol sorbitan fatty acid ester.

The use of disintegrators is also known from the patent literature. Thus, according to German patent application **DE 938 566**, acetyl salicylic acid is converted into granular form before compression, carefully but completely dried and then coated with highly disperse silica. The acetyl salicylic acid granules powdered with highly disperse silica may then be mixed with other tablet ingredients present in powder form or granular form and compressed to tablets. The barrier layer of highly disperse silica not only acts as an insulating layer and affords protection against unwanted reactions, it also contributes towards the rapid disintegration of the tablets, even after prolonged storage.

German patent application **DE 12 28 029** describes the production of tablets in which powder mixtures are mixed with cellulose powder and optionally highly disperse silica without preliminary granulation and the resulting mixture is compressed, preferably after grinding.

According to German patent application **DE 41 21 127**, a particularly effective auxiliary in the production of pharmaceutical tablets comprises cellulose particles with a coating material fixed to their surface. The auxiliary is used in fine-particle form, mean particle sizes below 200 μm being said to be particularly advantageous. Normally, therefore, traditional tablet disintegrators belonging to the first class of substances mentioned

are either mixed in very fine-particle form with the other tablet ingredients, which may be present as fine-particles or granules, before the compression step or the other tablet ingredients are coated or powdered with the tablet disintegrator.

5 According to the teaching of European patent **EP-B-0 523 099**, disintegrators known from the production of pharmaceuticals may also be used in detergents or cleaning compositions. The disintegrators mentioned include swellable layered silicates, such as bentonites, starch- and cellulose-based natural materials and derivatives thereof, alginates and the
10 like, potato starch, methyl cellulose and/or hydroxypropyl cellulose. These disintegrators may be mixed with the granules to be compressed or may even be incorporated therein.

 According to International patent application **WO-A-96/06156** also, it can be of advantage to incorporate disintegrators in detergent tablets. In
15 this case, too, typical disintegrators are said to include microcrystalline cellulose, sugars, such as sorbitol, and layered silicates, more particularly fine-particle and swellable layered silicates of the bentonite and smectite type. Effervescent substances, such as citric acid, bisulfate, bicarbonate, carbonate and percarbonate, are also mentioned as possible disintegration
20 aids.

 European patent applications **EP-A-0 466 485**, **EP-A-0 522 766**, **EP-A-0 711 827**, **EP-A-0 711 828** and **EP-A-0 716 144** describe the production of deterative tablets in which compacted particulate material with a particle size of 180 to 2000 μm is used. The resulting tablets may have both a
25 homogeneous structure and a heterogeneous structure. According to **EP-A-0 522 766**, at least those particles which contain surfactants and builders are coated with a solution or dispersion of a binder/disintegration aid, more particularly polyethylene glycol. Other binders/disintegration aids are - again - the already repeatedly described and known disintegrators, for
30 example starches and starch derivatives, commercially available cellulose

derivatives, such as crosslinked and modified cellulose, microcrystalline cellulose fibers, crosslinked polyvinyl pyrrolidones, layered silicates, etc. Weak acids, such as citric acid or tartaric acid which, in conjunction with carbonate-containing sources, lead to effervescent effects on contact with water and which, according to Römpp's definition, belong to the second class of disintegrators, may also be used as a coating material.

Particular reference is made to hitherto unpublished **DE 197 10 254** which describes disintegrators of which the particle size distribution (sieve analysis) is such that at most 1% by weight, preferably less, of dust is present and a total (including any dust present) of less than 10% by weight of the disintegrator granules are smaller than 0.2 mm. At least 90% by weight of the disintegrator granules advantageously have a particle size of at least 0.2 mm and at most 3 mm.

Disintegrators belonging to the group of organic acids, for example citric acid, or a citric acid/bicarbonate mixture and/or the celluloses and cellulose derivatives are also suitable. If a disintegrator is present in the shaped body, the dissolving time of the shaped body as a whole is preferably shorter than the duration of the main wash cycle of a conventional dishwashing machine, i.e. shorter than 40 mins., preferably shorter than 30 mins., more preferably shorter than 20 mins., and most preferably shorter than 10 mins.

In general, paraffins and/or microwaxes and/or the high molecular weight polyethylene glycols described in detail in the prior art are normally used as dissolution retarders. The use of mixtures mentioned in hitherto unpublished **DE 197 27 073**, of which the disclosure is hereby specifically included as part of the present disclosure, is particularly suitable for the purposes of the present application.

If the shaped body contains a component for retarding solubility, the dissolving time of the shaped body as a whole in cold (20°C) water - in one preferred embodiment - is longer than the preinse cycle of a commercially

available dishwashing machine, i.e. longer than 5 mins. and preferably longer than 10 mins.

General forms of presentation

5 The shaped body may assume any of the various three-dimensional forms which are listed inter alia in patents belonging to the prior art and in the standard works of the specialist literature (for example: **Riedel, "Die Tablette"**). The region in which the ingredient (I) is contained is also not confined to any particular three-dimensional form. However, a dimension
10 in one direction in space of more than 5 mm is preferred for practical reasons.

Specific forms of presentation

 In one preferred embodiment, the invention provides homogeneous
15 or heterogeneous shaped bodies of known construction.

 These include in particular cylindrical tablets which preferably have a diameter of 15 to 60 mm and, more particularly, 30 ± 10 mm. The height of these tablets is preferably from 5 to 30 mm and more preferably from 15 to 28 mm. Shaped bodies with a diameter of 32, 33, 34, 35, 36, 37, 38 and
20 39 mm have proved to be particularly favorable. In particular embodiments, the height is 24, 25, 26, 27 or 28 mm.

 However, square, rectangular, trapezoidal, oval and irregularly shaped bases may also be used with advantage. The edge lengths are preferably between 15 and 60 mm and more preferably 30 ± 10 mm.

25 The weight of the individual shaped bodies, more particularly tablets, is preferably 15 to 60 g and more preferably 20 to 40 g per shaped body or tablet. By contrast, the density of the shaped bodies or tablets normally assumes values above 1 kg/dm^3 and preferably in the range from 1.1 to 1.4 kg/dm^3 .

30 One or more shaped bodies, for example 2 to 4 shaped bodies,

more particularly tablets, may be used according to the particular application, the water hardness range and the nature of the soils to be removed. Other shaped bodies according to the invention may have even smaller diameters or dimensions, for example around 10 mm.

5 A homogeneous shaped body in the context of the present invention is understood to be one in which the ingredients of the shaped body - except for the region containing ingredient (I) - are homogeneously distributed in such a way that differences between ingredients and/or active substances are not visible to the naked eye. The granular structure of the
10 solids used is of course still discernible. In one preferred embodiment of the invention, only one other region (phase) is present besides the region containing the ingredient (I).

 Accordingly, heterogeneous shaped bodies in the context of the present invention are those in which the ingredients present besides the
15 ingredient (I) do not have to be homogeneously distributed. In one simple case, heterogeneous shaped bodies may be produced, for example, by giving the various ingredients different colors and/or by providing them with different fragrance components.

 Another case which, according to the invention, counts as a non-
20 homogeneous (heterogeneous) shaped body is an embodiment in which a shaped body comprising several layers (phases), i.e. at least two layers, besides the region containing the ingredient (I) is compressed. For example, these various layers may have different disintegration and dissolving rates and/or may contain different ingredients. This can result in
25 advantageous performance properties of the shaped bodies. If, for example, ingredients which adversely affect one another are present in the shaped bodies, they can be separated. If a particular sequence of cleaning conditions is to be created in a machine, one (or more) component(s) may be integrated in a (or the) faster disintegrating and/or faster dissolving layer
30 while the other component(s) is incorporated in a (or the) more slowly

dissolving layer, so that one component can act in advance or can have already reacted off by the time another component dissolves.

In one preferred embodiment of the invention, another two phases are present besides the region containing the ingredient (I). It is particularly favorable if the ratios by volume of the other two phases are
5 between 10:1 and 1:10, preferably between 5:1 and 1:5 and more preferably between 2:1 and 1:2.

Another particularly preferred embodiment is characterized in that three or more other phases are present.

10 The layered structure of the shaped bodies may be stack-like, in which case the inner layer(s) dissolve(s) at the edges of the shaped body before the outer layers have completely dissolved or disintegrated. However, the inner layer(s) may also be completely and/or partly surrounded by the layers situated further to the outside which prevents
15 and/or retards the early dissolution of constituents of the inner layer(s).

In another preferred embodiment of the invention, a tablet consists of at least three layers, i.e. two outer layers and at least one inner layer, at least one of the inner layer(s) containing a peroxy bleaching agent whereas, in the case of the stack-like tablet, the two outer layers and - in
20 the case of the jacket-like tablet - the outermost layers are free from peroxy bleaching agent. Peroxy bleaching agent and any bleach activators or bleach catalysts and/or enzymes present can also be separated from one another in one and the same tablet and/or shaped body. Embodiments such as these have the advantage that there is no danger of any unwanted
25 interactions in the shaped bodies according to the invention.

In another favorable and preferred embodiment of the present invention, one of the bleach components or the bleach component, more especially the chlorine component, is not made up in one phase with the perfume component. Another favorable and preferred embodiment of the
30 invention is characterized in that the silver protector component is not

made up with (any of) the bleach component(s). In another favorable and preferred embodiment of the invention, one of the components, or the component, for controlling solubility is made up together with the bleach activator. Another favorable and preferred embodiment of the invention is characterized in that one of the components, or the component, for controlling solubility is made up together with enzymes. Another favorable and preferred embodiment of the invention is characterized in that one of the components, or the component, for controlling solubility is made up together with the bleaching agent. Another favorable and preferred embodiment of the invention is characterized in that one of the components, or the component, for controlling solubility is made up together with the silver protector component. Another favorable and preferred embodiment of the invention is characterized in that one of the components, or the component, for controlling solubility is made up together with at least 50% by weight, preferably more than 70% by weight and, more preferably, more than 90% by weight of a surfactant or the surfactant mixture as a whole.

The shaped bodies according to the invention are produced by any of the methods normally used for producing shaped bodies.

In some embodiments, it has proved to be particularly favorable to introduce the region containing the ingredient(s) (I) in the form of a liquid (quite generally a solution and/or melt), optionally at elevated temperature, into a preformed recess of the surrounding ingredients and then to allow this region to solidify by applying conventional technical measures (quite generally drying and/or cooling). A viscosity of more than 1,500 mPas, preferably more than 2,000 mPas, more preferably between 2,000 and 15,000 mPas, most preferably between 2,500 and 7,000 mPas and, in a particularly advantageous embodiment, between 3,000 and 4,000 mPas, has proved to be particularly effective.

It has proved to be particularly effective to combine the active

ingredient (I) with fusible carrier materials. These include in principle any substances which solidify at or particularly above room temperature.

According to the invention, nonionic surfactants (Dehypon® LT 104, Dehypon® LS 54, Dehydol® LS 30, Lutensol® AT 80), polyethylene glycols
5 with various molecular weights (PEG 400, 12000), soaps (Lorol® C 16),
stearates (Cutina® GMS), and also thickened caustic soda and fusible
salts, such as sodium carbonate decahydrate, have proved to be
particularly suitable carrier materials. The advantages and disadvantages
of the individual ingredients can be reproduced by the expert in tests.

10 The use of mixtures mentioned in hitherto unpublished **DE 197 27 073**, of which the disclosure is hereby specifically included as part of the
present disclosure, is particularly suitable for the purposes of the present
application.

In other embodiments, parts of the shaped body and/or components
15 of the shaped body are treated by exposure to microwave radiation in order
positively to influence their strength, moisture content and solubility.

However, the constituents may also be simply compressed. In this
case, microtabs and shaped bodies smaller in their dimensions than the
final shaped body may be used in addition to powders and granules.

20

Ingredients

Particular ingredients (I) used in accordance with the invention are
substances which prevent the resoiling of surfaces and/or which facilitate
the separation of soil after a single application (so-called soil-release
25 compounds).

The soil-release compounds used in accordance with the invention
include any of the compounds known from the prior art. Cationic polymers
known, for example, from the following documents are particularly suitable:

According to **EP-A-0 167 382**, **EP-A-0 342 997** and **DE-OS 26 16**
30 **404**, cationic polymers are added to cleaning compositions to obtain

streak-free cleaning of the surfaces.

EP-A-0 167 382 describes liquid cleaning compositions which may contain cationic polymers as thickeners. Hydroxypropyl trimethyl ammonium guar; copolymers of aminoethyl methacrylate and acrylamide and
5 copolymers of dimethyl diallyl ammonium chloride and acrylamide are described as particularly suitable cationic polymers.

EP-A-0 342 997 describes multipurpose cleaners which may contain cationic polymers, polymers containing imino groups in particular being used.

10 DE-OS 26 16 404 describes cleaning compositions for glass containing cationic cellulose derivatives. The addition of the cationic cellulose derivatives to the compositions ensures better drainage of the water so that streak-free glass is obtained.

EP-A-0 467 472, for example, describes cleaning compositions for
15 hard surfaces which contain cationic homopolymers and/or copolymers as soil-release polymers. These polymers contain quaternized ammonium alkyl methacrylate groups as monomer units. These compounds are used to finish the surfaces in such a way that the soils can be removed more easily the next time the surfaces are cleaned.

20 Particular preference is attributed to the cationic polymers selected from cationic polymers of copolymers of such monomers as trialkyl ammonium alkyl (meth)acrylate or acrylamide; dialkyl diallyl diammonium salts; polymer-analog reaction products of ethers or esters of polysaccharides with lateral ammonium groups, more particularly guar, cel-
25 lulose and starch derivatives; polyadducts of ethylene oxide with ammonium groups; quaternary ethylene imine polymers and polyesters and polyamides containing quaternary lateral groups as soil-release compounds.

Natural polyuronic acids and related substances and also polyampholytes and hydrophobicized polyampholytes and mixtures of
30 these substances are also particularly preferred for the purposes of the

present invention.

According to the invention, between 0 and 5% by weight of enzymes, based on the composition as a whole, may be added to the detergent/cleaner to improve performance or guarantee the same quality of cleaning under milder conditions. The most commonly used enzymes include lipases, amylases, cellulases and proteases. Preferred proteases are, for example, BLAP® 140 (Biozym), Optimase® M-440 and Opticlean® M-250 (Solvay Enzymes); Maxacal® CX and Maxapem® or Esperase® (Gist Brocades) and even Savinase® (Novo). Particularly suitable celluloses and lipases are Celluzym® 0,7 T and Lipolase® 30 T (Novo Nordisk). Particularly suitable amylases are Duramyl® and Termamyl® 60 T and Termamyl® 90 T (Novo), Amylase-LT® (Solvay Enzymes) and Maxamyl® P5000 (Gist Brocades). Other enzymes may also be used.

15 Other ingredients

Particular ingredients (I) used in accordance with the invention are oxygen bleaching agents, preferably the alkali metal perborates and hydrates thereof and alkali metal percarbonates, sodium perborate (monohydrate or tetrahydrate) or sodium percarbonate and hydrates thereof preferably being used in accordance with the invention. The persulfates may also be used.

However, other typical oxygen bleaching agents are the organic peracids. Organic acids preferably used include above all the extremely effective phthalimidoperoxycaproic acid although, in principle, any other known peracids may also be used.

Particular ingredients (I) used in accordance with the invention are bleach activators. Known bleach activators are compounds which contain one or more N- or O-acyl groups, such as compounds from the class of anhydrides, esters, imides and acylated imidazoles or oximes. Examples are tetraacetyl ethylenediamine (TAED), tetraacetyl methylenediamine

(TAMD) and tetraacetyl hexylenediamine (TAHD) and also pentaacetyl glucose (PAG), 1,5-diacetyl-2,2-dioxohexahydro-1,3,5-triazine (DADHT) and isatoic anhydride (ISA).

The bleach activators used may be compounds which form aliphatic
5 peroxocarboxylic acids preferably containing 1 to 10 carbon atoms and, more particularly, 2 to 4 carbon atoms and/or optionally substituted perbenzoic acid under perhydrolysis conditions. Suitable bleach activators are substances which carry O- and/or N-acyl groups with the number of carbon atoms mentioned and/or optionally substituted benzoyl groups. Preferred
10 bleach activators are polyacylated alkylenediamines, more especially tetraacetyl ethylenediamine (TAED), acylated triazine derivatives, more especially 1,5-diacetyl-2,4-dioxohexahydro-1,3,5-triazine (DADHT), acylated glycolurils, more especially tetraacetyl glycoluril (TAGU), N-acyl imides, more especially N-nonanoyl succinimide (NOSI), acylated phenol
15 sulfonates, more particularly n-nonanoyl or isononanoyloxybenzenesulfonate (n- or iso-NOBS), carboxylic anhydrides, more especially phthalic anhydride, acylated polyhydric alcohols, more especially triacetin, ethylene glycol diacetate, 2,5-diacetoxy-2,5-dihydrofuran, n-methyl morpholinium acetonitrile methyl sulfate (MMA) and the enol esters known from German
20 patent applications **DE 196 16 693** and **DE 196 16 767** and also acetylated sorbitol and mannitol and the mixtures thereof (SORMAN) described in European patent application **EP 0 525 239**, acylated sugar derivatives, more especially pentaacetyl glucose (PAG), pentaacetyl fructose, tetraacetyl xylose and octaacetyl lactose and acetylated, optionally N-alkylated, glucamine and gluconolactone, and/or the N-acylated lactams,
25 for example N-benzoyl caprolactam, which are known from International patent applications **WO 94/27970**, **WO 94/28102**, **WO 94/28103**, **WO 95/00626**, **WO 95/14759** and **WO 95/17498**. The hydrophilically substituted acyl acetals known from German patent application **DE 196 16**
30 **769** and the acyl lactams described in German patent application **DE 196**

16 770 and in International patent application **WO 95/14075** are also preferably used. The combinations of conventional bleach activators known from German patent application **DE 44 43 177** may also be used. Bleach activators such as these are present in the usual quantities, preferably in quantities of 1% by weight to 10% by weight and more preferably in quantities of 2% by weight to 8% by weight, based on the detergent/cleaner as a whole.

In addition to or instead of the conventional bleach activators mentioned above, the sulfonimines known from European patents **EP 0 446 982** and **EP 0 453 003** and/or bleach-boosting transition metal salts or transition metal complexes may be present as so-called bleach catalysts. Suitable transition metal compounds include in particular the manganese-, iron-, cobalt-, ruthenium- or molybdenum-salen complexes known from German patent application **DE 195 29 905** and the N-analog compounds thereof known from German patent application **DE 196 20 267**, the manganese-, iron-, cobalt-, ruthenium- or molybdenum-carbonyl complexes known from German patent application **DE 195 36 082**, the manganese, iron, cobalt, ruthenium, molybdenum, titanium, vanadium and copper complexes with nitrogen-containing tripod ligands described in German patent application **DE 196 05 688**, the cobalt-, iron-, copper- and ruthenium-ammine complexes known from German patent application **DE 196 20 411**, the manganese, copper and cobalt complexes described in German patent application **DE 44 16 438**, the cobalt complexes described in European patent application **EP 0 272 030**, the manganese complexes known from European patent application **EP 0 693 550**, the manganese, iron, cobalt and copper complexes known from European patent **EP 0 392 592** and/or the manganese complexes described in European patent **EP 0 443 651** or in European patent applications **EP 0 458 397**, **EP 0 458 398**, **EP 0 549 271**, **EP 0 549 272**, **EP 0 544 490** and **EP 0 544 519**.

Combinations of bleach activators and transition metal bleach catalysts are

known, for example, from German patent application **DE 196 13 103** and International patent application **WO 95/27775**. Bleach activators from the group of polyacylated alkylenediamines, more particularly tetraacetyl ethylenediamine (TAED), N-acyl imides, more particularly N-nonanoyl succinimide (NOSI), acylated phenol sulfonates, more particularly n-nonanoyl or isononanoyloxybenzenesulfonate (n- or iso-NOBS), MMA are preferably used, preferably in quantities of up to 10% by weight, more preferably in quantities of 0.1% by weight to 8% by weight, most preferably in quantities of 2 to 8% by weight and, with particular advantage, in quantities of 2 to 6% by weight, based on the detergent/cleaner as a whole.

Bleach-boosting transition metal complexes, more particularly containing the central atoms Mn, Fe, Co, Cu, Mo, V, Ti and/or Ru, preferably selected from the group of manganese and/or cobalt salts and/or complexes, more preferably the cobalt (ammine) complexes, cobalt (acetate) complexes, cobalt (carbonyl) complexes, chlorides of cobalt or manganese and manganese sulfate, are also present in typical quantities, preferably in a quantity of up to 5% by weight, more preferably in a quantity of 0.0025% by weight to 1% by weight and most preferably in a quantity of 0.01% by weight to 0.25% by weight, based on the detergent/cleaner as a whole. In special cases, however, more bleach activator may even be used.

Dishwasher detergents according to the invention may contain corrosion inhibitors as an ingredient for protecting the tableware or the machine itself, silver protectors being particularly important for dishwashing machines. Known corrosion inhibitors, for example those described in **DE 43 25 922**, **DE 41 28 672** and **DE 43 38 724**, may be used. Above all, silver protectors selected from the group of triazoles, benzotriazoles, bisbenzotriazoles, aminotriazoles, alkylaminotriazoles and the transition metal salts or complexes may generally be used. Benzotriazole and/or alkylaminotriazole is/are particularly preferred. In addition, dishwashing

formulations often contain corrosion inhibitors containing active chlorine which are capable of distinctly reducing the corrosion of silver surfaces. According to the above documents, chlorine-free dishwashing detergents contain in particular oxygen- and nitrogen-containing organic redox-active compounds, such as dihydric and trihydric phenols, for example hydroquinone, pyrocatechol, hydroxyhydroquinone, gallic acid, phloroglucinol, pyrogallol and derivatives of these compounds. Salt-like and complex-like inorganic compounds, such as salts of the metals Mn, Ti, Zr, Hf, V, Co and Ce are also frequently used. Of these, the transition metal salts selected from the group of manganese and/or cobalt salts and/or complexes are preferred, cobalt(ammine) complexes, cobalt(acetate) complexes, cobalt(carbonyl) complexes, chlorides of cobalt or manganese and manganese sulfate being particularly preferred. Zinc compounds may also be used to prevent corrosion of tableware.

15 In the shaped bodies according to the invention, all ingredients (I) described above can also perform the function of other ingredients should the condition of the region according to the invention be fulfilled by other ingredients.

20 Water-soluble and water-insoluble builders may be used in the detergents/cleaners according to the invention, above all to bind calcium and magnesium. Water-soluble builders are preferred because, in general, they tend less to form insoluble residues on tableware and hard surfaces. Typical builders which may be present in quantities of 10 to 90% by weight, based on the composition as a whole, in accordance with the invention are 25 the low molecular weight polycarboxylic acids and salts thereof, the homopolymeric and copolymeric polycarboxylic acids and salts thereof, the carbonates, phosphates and silicates. Water-insoluble builders include the zeolites, which may also be used, and mixtures of the builders mentioned above.

30 Trisodium citrate and/or pentasodium tripolyphosphate and/or

sodium carbonate and/or sodium bicarbonate and/or gluconates and/or silicate builders from the class of disilicates and/or metasilicates are preferably used.

Alkali sources may be present as further ingredients. Alkali sources
5 are alkali metal hydroxides, alkali metal carbonates, alkali metal hydrogen carbonates, alkali metal sesquicarbonates, alkali metal silicates, alkali metal metasilicates and mixtures of the substances mentioned above. According to the invention, alkali metal carbonates, more particularly sodium carbonate, sodium hydrogen carbonate or sodium sesqui-
10 carbonate, are preferably used.

A builder system containing a mixture of tripolyphosphate and sodium carbonate is particularly preferred.

Another particularly preferred builder system contains a mixture of tripolyphosphate and sodium carbonate and sodium disilicate.

15 In principle, any surfactants may be used as surfactants. Nonionic surfactants, above all low-foaming nonionic surfactants, are preferred although other low-foaming surfactants may also be used. Alkoxyated alcohols, especially ethoxylated and/or propoxylated alcohols, alkyl polyglycosides, and alkyl polyglucamides are particularly preferred.

20 Alkoxyated alcohols are generally understood by the expert to be the reaction products of alkylene oxide, preferably ethylene oxide, with alcohols, the relatively long-chain alcohols (C_{10} to C_{18} , preferably between C_{12} and C_{16} such as, for example, C_{11} , C_{12} , C_{13} , C_{14} , C_{15} , C_{16} , C_{17} and C_{18} alcohols) being preferred for the purposes of the present invention. In
25 general, a complex mixture of addition products differing in their degree of ethoxylation is formed from n moles of ethylene oxide and 1 mole of alcohol, depending on the reaction conditions. Another embodiment is characterized by the use of mixtures of alkylene oxides, preferably a mixture of ethylene oxide and propylene oxide. If desired, "end-capped"
30 alcohol ethoxylates, which may also be used in accordance with the

invention, may also be obtained by etherification with short-chain alkyl groups, preferably butyl groups, in a concluding step. According to the invention, highly ethoxylated fatty alcohols or mixtures thereof with end-capped fatty alcohol ethoxylates are particularly preferred for the purposes
5 of the invention.

Alkyl polyglycosides are surfactants which may be obtained by reacting sugars and alcohols using the relevant methods of preparative organic chemistry, a mixture of monoalkylated oligomeric or polymeric sugars being obtained according to the particular method of production
10 used. Preferred alkyl polyglycosides are alkyl polyglucosides. In a particularly preferred embodiment, the alcohol is a long-chain fatty alcohol or a mixture of long-chain fatty alcohols while the degree of oligomerization of the sugars is between 1 and 10.

Fatty acid polyhydroxylamides (glucamides) are acylated reaction
15 products of the reductive amination of a sugar (glucose) with ammonia, long-chain fatty acids, long-chain fatty acid esters or long-chain fatty acid chlorides generally being used as acylating agents. Secondary amides are formed if methyl amine or ethyl amine is used instead of ammonia for the reduction process, as described for example in **SÖFW Journal**, 119,
20 (1993), 794-808. Carbon chain lengths of C₆ to C₁₂ are preferably used in the fatty acid moiety.

The regions of the shaped body may be colored. In one particularly preferred embodiment, one or more regions in the shaped body or all the regions in the shaped body are differently colored. In one special
25 embodiment of the shaped body, the color is red. In another special embodiment of the shaped body, the color is green. In another special embodiment of the shaped body, the color is yellow. In another special embodiment of the shaped body, the color is a mixture of different colors.

The storage stability of the shaped body is particularly important. In
30 one particularly preferred embodiment of the invention, the increase in

weight of the region containing the ingredient (I) during storage under normal domestic conditions, i.e. between 15 and 30°C and between 5 and 55% and preferably between 15 and 35% air humidity for a period of 30 days, preferably 60 days and more preferably 90 days is no more than 50%
5 by weight, preferably no more than 40% by weight, more preferably no more than 30% by weight, most preferably no more than 20% by weight and, in one particularly advantageous embodiment, no more than 10% by weight and, in the most favorable case, no more than 5% by weight.

In another particularly preferred embodiment of the invention, the
10 loss of active substance of the ingredient (I) during storage under normal domestic conditions, i.e. between 15 and 30°C and between 5 and 55% and preferably between 15 and 35% air humidity for a period of 30 days, preferably 60 days and more preferably 90 days is no more than 50% by weight, preferably no more than 40% by weight, more preferably no more
15 than 30% by weight, most preferably no more than 20% by weight and, in one particularly advantageous embodiment, no more than 10% by weight and, in the most favorable case, no more than 5% by weight.

In another particularly preferred embodiment of the invention, the absorption maximum of a 1% solution of a colored region or a colored
20 component during storage under normal domestic conditions, i.e. between 15 and 30°C and between 5 and 55% and preferably between 15 and 35% air humidity for a period of 30 days, preferably 60 days and more preferably 90 days is no more than 100 wave numbers, preferably no more than 50 wave numbers, more preferably no more than 30 wave numbers, most
25 preferably no more than 20 wave numbers and, in one particularly advantageous embodiment, no more than 10 wave numbers and, in the most favorable case, no more than 5 wave numbers.

Examples

5 The following formulations were mixed and compressed to shaped bodies. The ingredients shown in brackets are examples of the class of compounds in question, but may be replaced by other substances listed in the application. The starting formulations represent preferred ranges of the invention.

10 In the measurement of effectiveness, the particular ingredient (I) selected was transferred to the region according to the invention. In the other region (other regions), it was reduced in accordance with the invention or left out altogether before the studies were carried out.

Possible starting formulation (Ra) and test formulations (Va) (figures = % by weight):

Ingredient	Ra	Va1	Va2	Va3	Va4	Va5	Va6	Va7
Phosphate (sodium tripolyphosphate)	30-60	54.8	56.4	30.5	43	48	48	47
Sodium carbonate	5-25	15.9	13.2	21.5	7	20	12	24
Sodium disilicate	0-40	7.3	7.5	33.7	20	3	14	
Polymer (Sokalan CP5)	0-10	2.2	2.2			1		5
Sodium metasilicate	0-10			1.8				0
Sodium hydrogen carbonate	0-10				1		3	
Bleach (sodium perborate)	2-20	9.0	9.0	6.7	15	10	12	9
Bleach activator (TAED) (co-pentammine-Cl)	0-5	2.1	2.1		3	2	1	1.5
Enzyme (amylase) (Duramyl 60 T)	0-5	2.0	2.0	0.7	3	3	3	2.5
Enzyme (protease) (Blap 200 S)	0-5	1.8	1.8	1.0	2	2	3	2.5
Phosphonate or phosphonic acid	0-5	0.9	0.9		2			1.5
Nonionic surfactant (Dehypon LS 54)	0-5	2.0	2.0	2.0	2	4	3	1.5
Silver protector (benzotriazole)	0-3	0.3	0.3	0	0.3	0.2		0.5
Tabletting aid (polyethylene glycol)	0-10					6		
Paraffin (Mp.: 53°C)	0-10		2.0	1.0				3
Perfume	0-1	0.2	0.2	0.2	1	0.4	1	0.5
Dye	0-4	1.4	1.3	1.4	0.7	0.4		1.5

The builder system of the above starting formulation may also have the following composition (all other ingredients same as for a).

Ingredient	Rb	Vb1	Vb2	Vb3	Vb4	Vb5	Vb6	Vb7
Phosphate (sodium tripolyphosphate)	30-60	59	56	53	48	48	48	47
Carbonate (sodium carbonate)	5-25	21.2	20.3	22.7	17	20	16	24
Polymer (Sokalan CP5)	0-10		2	10	5	4		5
Sodium metasilicate	0-10			1.8				
Sodium hydrogen carbonate	0-10				1		3	

Examples of another starting formulation (R) according to the present invention and test formulations (V) (% by weight) are shown in the following Table:

Ingredient	Rc	Vc1	Vc2	Vc3	Vc4
Phosphate (sodium tripolyphosphate)	15-35	30	21	35	40
Carbonate (sodium carbonate)	25-55	39	50	40	35
Sodium hydrogen carbonate	10-20	14	19	15	12
Polymer (Sokalan CP5)	0-10	3	0		
Sodium disilicate	0-10				
Sodium disilicate	0-10	0	0	0	5
Bleach (sodium perborate)	2-10	7	8	5	5
Bleach activator (TAED)	0-5	1	0	1	1
Enzyme (amylase) (Duramyl 60 T)	0.4-2	1	0.5	0.5	0.5
Enzyme (protease) (Blap 200 S)	0.4-2	1	0.5	0.5	0.5
Phosphonate or phosphonic acid	0-5	0	0	0	0
Nonionic surfactant (Dehypon LS 54)	0-5	2	1	1	1
Silver protector (benzotriazole)	0-3	0.5	0	0.3	0
Paraffin (Mp.: 53°C)	0-10	0	0		0
Perfume	0-1	0	0	0.7	0
Dye	0-4	0.5	0	1	0

Further examples of a starting formulation (R) according to the present application and test formulations (V) (% by weight) are shown in the following Table:

Ingredient	Rd	Vd1	Vd2	Vd3	Vd4
Trisodium citrate	20-55	44	34	44	44
Sodium hydrogen carbonate	5-35	24	9	34	20
Carbonate (sodium carbonate)	0-10	7			
Polymer (Sokalan CP5)	0-10		6		
Sodium disilicate	0-25		20		
Bleach (sodium perborate)	0-22	9	16	17	
Bleach activator (TAED)	0-25	3	1	0	20
Enzyme (amylase) (Duramyl 60 T)	0.4-5	2.5	3	1	5
Enzyme (protease) (Blap 200 S)	0.4-5	2	3	1	5
Phosphonate or phosphonic acid	0-5	1.5	2.3		2
Nonionic surfactant (Dehypon LS 54)	0-5	2.5	1.7	2	1

Ingredient	Rd	Vd1	Vd2	Vd3	Vd4
Silver protector (benzotriazole)	0-3	0.5	0.5		
Paraffin (Mp.: 53°C)	0-10	3	1.5		
Perfume	0-1	0.9	0.5	1	1
Dye	0-4	0.1	1.5		2

Further examples of a starting formulation (R) according to the present application and test formulations (V) (% by weight) are shown in the following Table:

Ingredient	Re	Ve1	Ve2	Ve3	Ve4
Citrate (trisodium citrate)	20-55	32	36	38	24
Phosphate (trisodium polyphosphate)	20-55	26	23	24	24
Carbonate (sodium carbonate)	1-15	11	10	11	15
Polymer (Sokalan CP5)	0-10		5	1.5	6
Silicate (sodium disilicate)	0-25	3			6
Bleach (sodium perborate)	2-20	16	3	12	9
Bleach activator (TAED)	0-5	0.5	5	2	3
Enzyme (amylase) (Duramyl 60 T)	0-5		5	3	2
Enzyme (protease) (Blap 200 S)	0-5		5	0.5	2
Phosphonate or phosphonic acid	0-5	2	2		2
Tabletting aid (polyethylene glycol)	0-10	6	4	5	0.8
Nonionic surfactant (Dehypon LS 54)	0-5	2.5		2.8	2
Silver protector (benzotriazole)	0-3			0.2	0.2
Paraffin (Mp.: 53°C)	0-10				
Perfume	0-1	1	1		1
Dye	0-4	0	1		1

Further examples of a starting formulation (R) according to the present application and test formulations (V) (% by weight) are shown in the following Table:

Ingredient	Rf	Vf1	Vf2	Vf3	Vf4
Phosphate (trisodium polyphosphate)	40-60	49	49	49	50
Carbonate (sodium carbonate)	0-20		17	19	5

Ingredient	Rf	Vf1	Vf2	Vf3	Vf4
Polymer (Sokalan CP5)	0-15	2	4	1	6
Silicate (sodium disilicate)	0-30	24	6	5	10
Bleach (sodium perborate)	0-15	10	8	10	8
Bleach activator (TAED)	0-5	1.5	2	3	2
Enzyme (amylase) (Duramyl 60 T)	0-5	2	2	1.5	2.5
Enzyme (protease) (Blap 200 S)	0-5	3.9	2	1.5	2.5
Phosphonate or phosphonic acid	0-8	0.8	1	1	2
Tabletting aid (polyethylene glycol)	0-10	0.2	4	3.7	5
Nonionic surfactant (Dehypon LS 54)	0-8	1.5	2	4	3
Silver protector (benzotriazole)	0-5	0.1	0.5	0.3	0.5
Perfume	0-2	1	0.5	0.5	0.5
Dye	0-4	3	2	0.5	1

Further examples of a starting formulation (R) according to the present application and test formulations (V) (% by weight) are shown in the following Table:

Ingredient	Rg	Vg1	Vg2	Vg3	Vg4
Phosphate (trisodium polyphosphate)	30-60	55.7	59.6	46.5	47
Sodium disilicate	5-40	22.3	17.5	39.2	24
Polymer (Sokalan CP5)	0-10	2.2	2.2		5
Sodium metasilicate	0-10			1.8	
Sodium hydrogen carbonate	0-10				
Bleach (sodium perborate)	2-20	10	10	6.7	10
Bleach activator (co-catalyst)	0-2	1.1	1.1		0.5
Enzyme (amylase) (Duramyl 60 T)	0-5	2.0	2.0	0.7	2.5
Enzyme (protease) (Blap 200 S)	0-5	1.8	1.8	1.0	2.5
Phosphonate or phosphonic acid	0-5	0.9	0.9		1.5
Nonionic surfactant (Dehypon LS 54)	0-5	2.0	2.0	2.0	1.5
Silver protector (benzotriazole)	0-3	0.3	0.3	0	0.5
Tabletting aid (polyethylene glycol)	0-10				
Paraffin (Mp.: 53°C)	0-10		2.0	1.0	3
Perfume	0-1	0.2	0.2	0.2	0.5
Dye	0-4	1.4	1.3	1.4	1.5

Further examples of a starting formulation (R) according to the present application and test formulations (V) (% by weight) are shown in the following Table:

Ingredient	Rh	Vh1	Vh2	Vh3	Vh4	Vh5	Vh6	Vh7
Phosphate (sodium tripolyphosphate)	30-60	40	40	40	45	45	50	35
Sodium metasilicate (5-hydrate)	5-45	45	10	40	10	30	20	35
Sodium metasilicate (0-hydrate)	5-40	10	10	8	10	8	5	10
Sodium disilicate	0-40	0	30	0	15	10	5	0
Paraffin oil	0-10	4	5	5	5	6	5	0
Bleach (trichloroisocyanuric acid)	0.5-10	1	2	2	3	1	5	5
Enzyme (amylase) (Duramyl 60 T)	0-5			1	2		2	2
Enzyme (protease) (Blap 200 S)	0-5			1	2		2	2
Nonionic surfactant (Dehypon LS 54)	0-5		2	2	2		2	3
Tabletting aid (polyethylene glycol)	0-10		1		4		2	4
Perfume	0-1		0.3	0.3	1		1	0.5
Dye	0-4		0.7	0.7	1		1	2.5

Another starting formulation (R) (% by weight) and test formulations (V) were tested:

Ingredient	Ri1	Vi1	Vi2	Vi3
Sodium carbonate	35-55	43.5	45	55
Sodium hydrogen carbonate	15-35	15	25	34
Polymer (Sokalan CP5)	3-10	5	3	0
Sodium metasilicate	0-10	3	2	0
Bleach (sodium perborate)	5-12	6	10	8
Bleach activator (TAED)	0-5	1	2	0
Enzyme (amylase) (Duramyl 60 T)	0.4-2	1	1	0.5
Enzyme (protease) (Blap 200 S)	0.4-2	1	1	0.5
Phosphonate or phosphonic acid	0-5	0	2	0
Nonionic surfactant (Dehypon LS 54)	0-5	1	2	1
Tabletting aid (polyethylene glycol)	0-5		3	
Silver protector (benzotriazole)	0-3	0.5	1	0
Paraffin (Mp.: 53°C)	0-10	0	2	0
Perfume	0-1	0	1	0
Dye	0-4	1	1	1

Should a shaped body according to the invention contain several phases, each formulation may also be the constituents of an individual phase, in other words the percentages are then not based - as usual - on the formulation as a whole, but rather on the composition of individual phase.

Each of the formulations may of course also contain other ingredients typical of MDWM (for example fillers, preservatives and the like) in extremely small quantities, in which case the percentage contents of the other ingredients should be varied accordingly.

The production of the shaped bodies according to the invention involves steps which are known to the expert in this field in another connection. A preferred embodiment of the shaped body according to

claim 1 contains a recess in a tablet, this recess containing the chlorine bleach ingredient (I). Production is preferably carried out by forming a depression in a shaped body and filling the depression thus formed. The depression may be formed in a Korsch rotary press. In the present case, a
5 Fette tablet press was used. A round tablet die (26 x 36 mm) was selected and, using a punch, a 5 mm deep recess was formed in one side, the base of the recess being selected so that a volume of 1 ml could be poured in, the surface of the tablet thereafter becoming smooth again.

A liquid mixture of paraffin and an active substance (I) was then
10 poured in and allowed to cool. The cooling process may be assisted by conventional methods and equipment.

As already mentioned, the ingredient (I) was quantitatively reduced beforehand in accordance with the invention in the corresponding formulation surrounding the filled region and, in the present Examples and
15 studies, was left out in the surrounding regions.

In comparisons with homogeneous tablets (where ingredient (I) does not correspond to the invention) and a few commercially available tablets, the tablets according to the invention proved to be superior.

The particular properties of the detergents/cleaners according to the
20 invention were tested by comparison with known detergents/cleaners using using the removal of tea stains as an example. The particular properties of the detergents/cleaners according to the invention were tested by comparison with known detergents/cleaners using using the removal of enzyme-relevant stains as an example.

25 In addition, the particular properties of the detergents/cleaners according to the invention were tested by technically untrained people under simulated practical conditions in comparison with homogeneous tablets. Apart from dosing instructions and general safety instructions, the handling of the detergents/cleaners by the people in question in
30 commercially available dishwashing machines was not influenced in any

way. This comparison test produced particularly good results. The cleaning results were evaluated as particularly favorable.

In a test, samples according to starting formulation Ra which contained active chlorine (dichlorocyanurate) as bleaching component on the one hand in homogeneous distribution in the tablet and, on the other hand, in accordance with claim 1, so that the active chlorine was largely separated from the other (particularly enzyme-containing) components, were used. Storage stability after 2 weeks was distinctly poorer in the case of the homogeneous tablet so it was not possible to obtain a satisfactory cleaning effect against milk and starch-relevant stains. This applies to the entirely normal making-up of the bleach component in region (I) which, in the present case, represented a recess. The effect is even clearer where a mixture of chlorine component and paraffin is used in a standard program at 55°C. The cleaning results were better even immediately after production. The effect improved with storage. Tea, milk, burnt-on meat, burnt-in meat, egg and starch mixtures in particular were removed far more effectively than by tablets without a region (I), i.e. with a homogeneously distributed component (I), and commercially available dishwasher tablets.

The effect was discernible even with paraffins having a melting point of 40 to 44°C. The effect is more pronounced with paraffins having a melting point of 46 to 48°C. The cleaning effect is particularly pronounced with paraffins having a melting point of 57 to 60°C.

The results clearly show that the tablets according to the invention are superior to conventional dishwasher tablets.

CLAIMS

1. A shaped body containing builders, alkali sources, bleaching agents, enzymes and surfactants, characterized in that more than 80% by weight of a mixture of active substance (I) in the form of a bleaching agent from the group of chlorine bleaching agents and a fusible component for controlling solubility selected from the group of surfactants, paraffins, microwaxes and relatively high molecular weight polyethylene glycols is contained in a region of no greater than 40% by volume.
2. A shaped body as claimed in claim 1, characterized in that a region of 1 to 30% by volume, preferably between 2 and 25% by volume and more preferably between 3 and 20% by volume of the shaped body contains more than 90% by weight, preferably more than 95% by weight and more preferably 100% by weight of an ingredient (I) present.
3. A shaped body as claimed in claim 1, characterized in that the ingredient (I) is a mixture of bleaching agent from the group of chlorine bleaching agents and another ingredient from the group consisting of bleach activator and/or silver protector and/or enzyme and/or a component for controlling solubility and/or other ingredients.
4. A shaped body as claimed in any of the preceding claims, characterized in that at least one bleaching agent selected from the group of trichlorocyanuric acids, dichloro- or monochlorocyanurates, hypochlorites and other typical chlorine-containing bleaching agents is present.
5. A shaped body as claimed in any of the preceding claims, characterized in that the fusible component is a paraffin.
6. A shaped body as claimed in claim 5, characterized in that the paraffin has a melting point of more than 40-44°C, preferably more than 46-48°C and more preferably between 57 and 60°C.
7. A shaped body as claimed in any of the preceding claims, characterized in that one or more builders from the group of gluconates,

low molecular weight polycarboxylic acids and salts thereof, homopolymeric and copolymeric polycarboxylic acids and salts thereof, carbonates, phosphates and silicates, preferably trisodium citrate and/or pentasodium tripolyphosphate and/or sodium carbonate and/or sodium bicarbonate and/or gluconates and/or silicate builders from the class of disilicates and/or metasilicates is/are also used.

8. A shaped body as claimed in any of the preceding claims, characterized in that it contains a mixture of tripolyphosphate and sodium carbonate as builder system.

10 9. A shaped body as claimed in any of the preceding claims, characterized in that it contains a mixture of tripolyphosphate and sodium carbonate and sodium disilicate as builder system.

10. A shaped body as claimed in any of the preceding claims, characterized in that the surfactants are selected from the group of nonionic surfactants, preferably low-foaming nonionic surfactants, more preferably alkoxylated and/or propoxylated fatty alcohols, alkyl polyglycosides, alkyl polyglucamides and/or mixtures of the surfactants mentioned.

11. A shaped body as claimed in any of the preceding claims, characterized in that at least one silver protector selected from the group of triazoles, benzotriazoles, bis-benzotriazoles, aminotriazoles, alkylaminotriazoles and transition metal salts or complexes is additionally present.

12. A shaped body as claimed in claim 11, characterized in that benzotriazole and/or alkyl aminotriazole is/are present as silver protector(s).

13. A shaped body as claimed in claim 11, characterized in that the transition metal salts are selected from the group of manganese and/or cobalt salts and/or complexes, preferably cobalt (ammine) complexes, cobalt (acetate) complexes, cobalt (carbonyl) complexes, chlorides of cobalt or manganese and manganese sulfate.

14. A shaped body as claimed in any of the preceding claims, characterized in that the component and/or components for controlling solubility are selected from the group of paraffins, microwaxes and high molecular weight polyethylene glycols.
- 5 15. A tablet as claimed in claim 14, characterized in that the dissolving time of the tablet as a whole is longer than the duration of the prerinse cycle of a conventional dishwasher.
16. A shaped body as claimed in any of claims 1 to 13, characterized in that the component and/or components for controlling solubility are
10 selected from the group of organic acids, for example citric acid, or a citric acid/bicarbonate mixture and/or celluloses and cellulose derivatives.
17. A tablet as claimed in claim 16, characterized in that the dissolving time of the tablet as a whole is shorter than the duration of the main wash cycle of a conventional dishwasher.
- 15 18. A shaped body as claimed in any of the preceding claims, characterized in that only one other region or one other phase is present.
19. A shaped body as claimed in any of claims 1 to 17, characterized in that two other regions or phases are present.
20. A shaped body as claimed in claim 19, characterized in that the
20 ratios by volume of the two other phases are between 10:1 and 1:10, preferably between 5:1 and 1:5 and more preferably between 2:1 and 1:2.
21. A shaped body as claimed in any of claims 1 to 17, characterized in that three or more other phases are present.
22. A shaped body as claimed in any of the preceding claims,
25 characterized in that one of the bleaching components or the bleaching component, more particularly the chlorine component, is not made up in one phase with the perfume component.
23. A shaped body as claimed in any of the preceding claims, characterized in that the silver protector is not made up with (any of) the
30 bleaching component(s).

24. A shaped body as claimed in any of the preceding claims, characterized in that one of the components, or the component, for controlling solubility is made up together with the bleach activator.

5 25. A shaped body as claimed in any of the preceding claims, characterized in that one of the components, or the component, for controlling solubility is made up together with the enzymes.

26. A shaped body as claimed in any of the preceding claims, characterized in that one of the components, or the component, for controlling solubility is made up together with the bleaching agent.

10 27. A shaped body as claimed in any of the preceding claims, characterized in that one of the components, or the component, for controlling solubility is made up together with the silver protector component.

15 28. A shaped body as claimed in any of the preceding claims, characterized in that one of the components, or the component, for controlling solubility is made up together with at least 50% by weight, preferably with more than 70% by weight and, more preferably, with more than 90% by weight of a surfactant or the mixture of surfactants as a whole.

20 29. A shaped body as claimed in any of the preceding claims, characterized in that one or more or all of the regions of the shaped body are differently colored.

30. A shaped body as claimed in any of the preceding claims, characterized in that one color is red.

25 31. A shaped body as claimed in any of the preceding claims, characterized in that one color is green.

32. A shaped body as claimed in any of the preceding claims, characterized in that one color is yellow.

33. A shaped body as claimed in any of the preceding claims, characterized in that the color is a mixture of colors.